



Arab Academy for Science & Technology & Maritime Transport

(AASTMT – Cairo Branch)

College of Engineering & technology

Electronics & Communication Engineering Department

Course : Solid State Electronics
Course Code : EC210
Instructors : Dr. Mostafa Fedawy
TA's : Eng. Nihal Sameh, Eng. Mahmoud Sadat

Sheet # 5

Schrodinger Equation

Physical Constants:

Charge of electron (e) = 1.6×10^{-19} C

Mass of proton (m_p) = 1.672×10^{-27} kg

Speed of light (C) = 3×10^8 m/s

1 eV = 1.6×10^{-19} J

Mass of electron (m_e) = 9.1×10^{-31} kg

Plank's constant (h) = 6.63×10^{-34} J.s

Put (T) for the true statement or (F) for the false statement:

1. If an electron with energy 5 eV impinges a semi-infinite potential step of height 3 eV, the probability that the electron is reflected back is zero.
2. If an electron with energy 3 eV impinges a semi-infinite potential step of height 5 eV, the probability that the electron is transmitted through the step is zero.
3. The tunneling probability through a potential barrier of finite width increases with the width of the barrier.
4. The tunneling probability through a potential barrier of finite width increases with the height of the barrier.
5. The tunneling probability through a potential barrier of finite width increases with the energy of the particle.

Choose the correct answer justifying your choice:

1. If an electron has a wave function of the form $A \exp[ikx]$, the probability of finding the electron at $x = 5$ is
 - (a) $A \exp[i 5k]$
 - (b) $A \exp[i 10k]$
 - (c) $A^2 \exp[i 10k]$
 - (d) A^2
 - (e) $A^2 \exp[i 5k]$
2. If an electron with energy 5 eV impinges a semi-infinite potential step of height 3 eV, the energy of the electron in the higher potential region will
 - (a) Decrease
 - (b) increase
 - (c) remain the same
 - (d) be zero
 - (e) Be negative


3. If an electron with energy 5 eV impinges a semi-infinite potential step of height 3 eV, the angular wave number of the electron in the higher potential region will
 - (a) Decrease
 - (b) increase
 - (c) Remain the same
 - (d) be zero
 - (e) Be negative

4. If an electron with energy 5 eV impinges a semi-infinite potential step of height 3 eV, the velocity of the electron in the higher potential region will
 - (a) Decrease
 - (b) increase
 - (c) Remain the same
 - (d) be zero
 - (e) Be negative

5. If an electron with energy 2 eV impinges a semi-infinite potential step of height 6 eV, the ratio of the probability of finding the electron in the higher potential region at a distance 2.25 \AA from the edge of the step to that at the edge of the step is approximately
 - (a) 0.1
 - (b) 10
 - (c) 10^{-3}
 - (d) 0.01
 - (e) Zero

Solve the following Problems:

1. An electron with energy $E = 2 \text{ eV}$ moves towards a potential step (shown in the figure) of height $V_o = 6 \text{ eV}$. Given $\psi_1 = Ae^{ikx} + Be^{-ikx}$, and $\psi_2 = Ce^{-\alpha x}$
 - (a) Calculate k and α .
 - (b) What is the reflection coefficient R ?
 - (c) What is the transmission coefficient T ?



2. An electron with a total energy E moves in a one dimensional region 1. At $x = 0$, there is a potential energy step of height V_o (as shown in the figure). Given $\psi_1 = Ae^{iax} + Be^{-iax}$, and $\psi_2 = Ce^{-x/b} + De^{x/b}$
 - (a) Find the real numbers a and b (in terms of E and V_o).
 - (b) Is the total energy of the electron less or greater than V_o ? why?
 - (c) Knowing that the wave function and its space derivative are single valued, continuous and finite find B, C, D in terms of A .

